arXiv:1212.2165v1 [math.CA] 7 Dec 2012

INEQUALITIES OF OSTROWSKI'S TYPE FOR $m-$ AND $(\alpha, m)-$ LOGARITHMICALLY CONVEX FUNCTIONS VIA RIEMANN-LIOUVILLE FRACTIONAL INTEGRALS

AHMET OCAK AKDEMIR

Abstract. In this paper, we establish some new Ostrowski's type inequalities for $m-$ and $(\alpha, m)-$ logarithmically convex functions by using the Riemann-Liouville fractional integrals.

1. INTRODUCTION

Let $f : I \subset [0, \infty] \to \mathbb{R}$ be a differentiable mapping on I°, the interior of the interval I, such that $f' \in L[a, b]$ where $a, b \in I$ with $a < b$. If $|f'(x)| \leq M$, then the following inequality holds (see [7]).

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(u) du \right| \leq \frac{M}{b-a} \left[\frac{(x-a)^2 + (b-x)^2}{2} \right] \tag{1.1}$$

This inequality is well known in the literature as the Ostrowski inequality. For some results which generalize, improve and extend the inequality (1.1) see ([7, 8, 9, 10, 11, 18]) and the references therein.

Let us recall some known definitions and results which we will use in this paper. The function $f : [a, b] \to \mathbb{R}$, is said to be convex, if we have

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

for all $x, y \in [a, b]$ and $t \in [0, 1]$. We can define starshaped functions on $[0, b]$ which satisfy the condition

$$f(tx) \leq tf(x)$$

for $t \in [0, 1]$.

The concept of $m-$convexity has been introduced by Toader in [5], an intermediate between the ordinary convexity and starshaped property, as following:

Definition 1. *The function* $f : [0, b] \to \mathbb{R}$, $b > 0$, *is said to be* $m-$*convex, where* $m \in [0, 1]$, *if we have*

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y)$$

for all $x, y \in [0, b]$ *and* $t \in [0, 1]$. *We say that* f *is* $m-$*concave if* $-f$ *is* $m-$*convex.*

In [4], Miheşan gave definition of $(\alpha, m)-$convexity as following;

2000 *Mathematics Subject Classification.* 26D15.

Key words and phrases. Ostrowski's inequality, $m-$ logarithmically convex functions, $(\alpha, m)-$ logarithmically convex.

Definition 2. *The function* $f:[0,b]\to\mathbb{R}$, $b>0$ *is said to be* $(\alpha,m)-$*convex, where* $(\alpha,m)\in[0,1]^2$, *if we have*

$$f(tx+m(1-t)y)\leq t^{\alpha}f(x)+m(1-t^{\alpha})f(y)$$

for all $x,y\in[0,b]$ *and* $t\in[0,1]$.

Denote by $K_m^{\alpha}(b)$ the class of all $(\alpha,m)-$convex functions on $[0,b]$ for which $f(0)\leq 0$. If we choose $(\alpha,m)=(1,m)$, it can be easily seen that $(\alpha,m)-$convexity reduces to $m-$convexity and for $(\alpha,m)=(1,1)$, we have ordinary convex functions on $[0,b]$. For the recent results based on the above definitions see the papers [2]-[9].

Definition 3. *([1]) A function* $f:[0,b]\to(0,\infty)$ *is said to be* $m-$*logarithmically convex if the inequality*

$$f\left(tx+m\left(1-t\right)y\right)\leq\left[f\left(x\right)\right]^{t}\left[f\left(y\right)\right]^{m(1-t)} \tag{1.2}$$

holds for all $x,y\in[0,b]$, $m\in(0,1]$, *and* $t\in[0,1]$.

Obviously, if putting $m=1$ in Definition 3, then f is just the ordinary logarithmically convex function on $[0,b]$.

Definition 4. *([1]) A function* $f:[0,b]\to(0,\infty)$ *is said to be* $(\alpha,m)-$*logarithmically convex if*

$$f\left(tx+m\left(1-t\right)y\right)\leq\left[f\left(x\right)\right]^{t^{\alpha}}\left[f\left(y\right)\right]^{m(1-t^{\alpha})} \tag{1.3}$$

holds for all $x,y\in[0,b]$, $(\alpha,m)\in(0,1]\times(0,1]$, *and* $t\in[0,1]$.

Clearly, when taking $\alpha=1$ in Definition 4, then f becomes the standard m-logarithmically convex function on $[0,b]$.

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 5. *Let* $f\in L_1[a,b]$. *The Riemann-Liouville integrals* $J_{a^+}^{\mu}f$ *and* $J_{b^-}^{\mu}f$ *of order* $\mu>0$ *with* $a\geq 0$ *are defined by*

$$J_{a^+}^{\mu}f\left(x\right)=\frac{1}{\Gamma(\mu)}\int_a^x\left(x-t\right)^{\mu-1}f(t)dt,\quad x>a$$

and

$$J_{b^-}^{\mu}f\left(x\right)=\frac{1}{\Gamma(\mu)}\int_x^b\left(t-x\right)^{\mu-1}f(t)dt,\quad x<b$$

respectively where $\Gamma(\mu)=\int_0^{\infty}e^{-t}u^{\mu-1}du$. *Here is* $J_{a^+}^{0}f(x)=J_{b^-}^{0}f(x)=f(x)$.

In the case of $\mu=1$, the fractional integral reduces to the classical integral. For some recent results connected with fractional integral inequalities see [11]-[18].

The aim of this study is to establish some Ostrowski type inequalities for the class of functions whose derivatives in absolute value are $m-$ and $(\alpha,m)-$ geometrically convex functions via Riemann-Liouville fractional integral.

2. THE NEW RESULTS

In order to prove our results, we need the following lemma that has been obtained in [11]:

Lemma 1. *([11]) Let $f:[a,b]\to\mathbb{R}$ be a differentiable mapping on (a,b) with $a<b$. If $f'\in L[a,b]$, then for all $x\in[a,b]$ and $\mu>0$ we have:*

$$\begin{aligned}&\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\\ =\;&\frac{(x-a)^{\mu+1}}{b-a}\int_0^1 t^{\mu}f'(tx+(1-t)a)\,dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1 t^{\mu}f'(tx+(1-t)b)\,dt\end{aligned}$$

where $\Gamma(\mu)=\int\limits_0^{\infty}e^{-t}u^{\mu-1}du$.

Theorem 1. *Let $f:[0,\infty)\to(0,\infty)$ be differentiable mapping with $a,b\in[0,\infty)$ such that $a<b$. If $|f'(x)|$ is $(\alpha,m)-$logarithmically convex function with $|f'(x)|\leq M$, $f'\in L[a,b]$, $(\alpha,m)\in(0,1]\times(0,1]$ and $\mu>0$, then the following inequality for fractional integrals holds:*

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\;&\left[\frac{1}{2\mu+1}+K_1(\alpha,m,t)\right]\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{2(b-a)}\right]\end{aligned}\tag{2.1}$$

where

$$K_1(\alpha,m,t)=\begin{cases}\frac{M^{2m}\left(M^{2\alpha-2\alpha m}-1\right)}{(2\alpha-2\alpha m)\ln M} & ,M<1\\ 1 & ,M=1\end{cases}.$$

Proof. By Lemma 1 and since $|f'|$ is $(\alpha,m)-$logarithmically convex, we can write

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\;&\frac{(x-a)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(tx+(1-t)a)\right|dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(tx+(1-t)b)\right|dt\\ \leq\;&\frac{(x-a)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(x)\right|^{t^{\alpha}}\left|f'\left(\frac{a}{m}\right)\right|^{m(1-t^{\alpha})}dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(x)\right|^{t^{\alpha}}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-t^{\alpha})}dt\\ \leq\;&\frac{(x-a)^{\mu+1}}{b-a}\int_0^1 t^{\mu}M^{m+t^{\alpha}(1-m)}dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1 t^{\mu}M^{m+t^{\alpha}(1-m)}dt.\end{aligned}$$

By using the elemantery inequality $cd\leq\frac{c^2+d^2}{2}$, we have

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\;&\frac{(x-a)^{\mu+1}}{b-a}\int_0^1\frac{t^{2\mu}+M^{2(m+t^{\alpha}(1-m))}}{2}dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1\frac{t^{2\mu}+M^{2(m+t^{\alpha}(1-m))}}{2}dt\\ =\;&\left[\frac{1}{2\mu+1}+\int_0^1 M^{2(m+t^{\alpha}(1-m))}dt\right]\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{2(b-a)}\right].\end{aligned}\tag{2.2}$$

If we choose $M=1$, then

$$\int_0^1 M^{2(m+t^{\alpha}(1-m))}dt=1.$$

If $M<1$, then $M^{2(m+t^{\alpha}(1-m))}\leq M^{2(m+\alpha t(1-m))}$, thus

$$\int_0^1 M^{2(m+\alpha t(1-m))}dt=\frac{M^{2m}\left(M^{2\alpha-2\alpha m}-1\right)}{(2\alpha-2\alpha m)\ln M}$$

Which completes the proof. □

Corollary 1. *Let $f:[0,\infty)\to(0,\infty)$ be differentiable mapping with $a,b\in[0,\infty)$ such that $a<b$. If $|f'(x)|$ is $m-$logarithmically convex function with $|f'(x)|\leq M, f'\in L[a,b]$, $m\in(0,1]$ and $\mu>0$, then the following inequality for fractional integrals holds:*

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ &\leq\left[\frac{1}{2\mu+1}+\frac{M^2-M^{2m}}{2\ln M-2m\ln M}\right]\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{2(b-a)}\right].\end{aligned}\tag{2.3}$$

Proof. If we take $\alpha=1$ in (2.1), we get the required result. □

Corollary 2. *Let $f:[0,\infty)\to(0,\infty)$ be differentiable mapping with $a,b\in[0,\infty)$ such that $a<b$. If $|f'(x)|$ is logarithmically convex function with $|f'(x)|\leq M, f'\in L[a,b]$ and $\mu>0$, then the following inequality for fractional integrals holds:*

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ &\leq\left[\frac{1}{2\mu+1}+M^2\right]\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{2(b-a)}\right].\end{aligned}\tag{2.4}$$

Proof. If we take $\alpha=m=1$ in (2.2), we get the required result. □

Corollary 3. *Let $f:[0,\infty)\to(0,\infty)$ be differentiable mapping with $a,b\in[0,\infty)$ such that $a<b$. If $|f'(x)|$ is logarithmically convex function with $|f'(x)|\leq M$ and $f'\in L[a,b]$, then the following inequality holds:*

$$\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right|\leq\left[\frac{1}{3}+M^2\right]\left[\frac{(x-a)^2+(b-x)^2}{2(b-a)}\right].$$

Proof. If we choose $\mu=1$ in (2.4), we get the required result. □

Theorem 2. *Let $f:[0,\infty)\to(0,\infty)$ be differentiable mapping with $a,b\in[0,\infty)$ such that $a<b$. If $|f'(x)|^q$ is $(\alpha,m)-$logarithmically convex function with $|f'(x)|^q\leq M, f'\in L[a,b]$, $(\alpha,m)\in(0,1]\times(0,1]$ and $\mu>0$, then the following inequality for fractional integrals holds:*

(2.5)

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ &\leq\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}(K_2(\alpha,m,t))^{\frac{1}{q}}\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{(b-a)}\right]\end{aligned}$$

where $q>1$, $0\leq p\leq q$ *and*

$$K_2(\alpha,m,t)=\begin{cases}\frac{M^m\left(\Gamma(\mu p+1)-\Gamma\left(\mu p+1,\ln M^{\alpha(m-1)}\right)\right)}{\left(\ln M^{\alpha(m-1)}\right)^{\mu p+1}} & ,M<1\\ \frac{1}{\mu p+1} & ,M=1\end{cases}.$$

Proof. From Lemma 1 and by using the properties of modulus, we have

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\ &\frac{(x-a)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(tx+(1-t)a)\right|dt+\frac{(b-x)^{\mu+1}}{b-a}\int_0^1 t^{\mu}\left|f'(tx+(1-t)b)\right|dt.\end{aligned}$$

By applying the Hölder inequality for $q>1$, $0\leq p\leq q$, we get

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\ &\frac{(x-a)^{\mu+1}}{b-a}\left[\left(\int_0^1 t^{\mu\left(\frac{q-p}{q-1}\right)}dt\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}\left|f'(tx+(1-t)a)\right|^q dt\right)^{\frac{1}{q}}\right.\\ &\left.+\frac{(b-x)^{\mu+1}}{b-a}\left(\int_0^1 t^{\mu\left(\frac{q-p}{q-1}\right)}dt\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}\left|f'(tx+(1-t)b)\right|^q dt\right)^{\frac{1}{q}}\right].\end{aligned}$$

It is easy to see that

$$\int_0^1 t^{\mu\left(\frac{q-p}{q-1}\right)}dt=\frac{q-1}{\mu(q-p)+q-1}.$$

Hence, by (α,m) −logarithmically convexity of $|f'|^q$, we have

$$\begin{aligned}&\left|\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right|\\ \leq\ &\frac{(x-a)^{\mu+1}}{b-a}\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}\left|f'(x)\right|^{t^{\alpha}}\left|f'\left(\frac{a}{m}\right)\right|^{m(1-t^{\alpha})}dt\right)^{\frac{1}{q}}\\ &+\frac{(b-x)^{\mu+1}}{b-a}\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}t^{\mu}\left|f'(x)\right|^{t^{\alpha}}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-t^{\alpha})}dt\right)^{\frac{1}{q}}\\ =\ &\frac{(x-a)^{\mu+1}}{b-a}\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}M^{m+t^{\alpha}(1-m)}dt\right)^{\frac{1}{q}}\\ &+\frac{(b-x)^{\mu+1}}{b-a}\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(\int_0^1 t^{\mu p}M^{m+t^{\alpha}(1-m)}dt\right)^{\frac{1}{q}}.\end{aligned}\tag{2.6}$$

If we choose $M=1$, then

$$\int_0^1 t^{\mu p}dt=\frac{1}{\mu p+1}.$$

If $M<1$, then $M^{m+t^{\alpha}(1-m)}\leq M^{m+\alpha t(1-m)}$, thus

$$\int_0^1 t^{\mu p}M^{m+\alpha t(1-m)}dt=\frac{M^{m}\left(\Gamma\left(\mu p+1\right)-\Gamma\left(\mu p+1,\ln M^{\alpha(m-1)}\right)\right)}{\left(\ln M^{\alpha(m-1)}\right)^{\mu p+1}}$$

Which completes the proof. □

Corollary 4. *Let* $f:[0,\infty)\rightarrow(0,\infty)$ *be differentiable mapping with* $a,b\in[0,\infty)$ *such that* $a<b$. *If* $\left\vert f'(x)\right\vert^{q}$ *is* $m-$*logarithmically convex function with* $\left\vert f'(x)\right\vert^{q}\leq M, f'\in L[a,b]$, $m\in(0,1]$ *and* $\mu>0$, *then the following inequality for fractional integrals holds:*

$$\begin{aligned}&\left\vert\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right\vert\\ \leq&\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(K_2(1,m,t)\right)^{\frac{1}{q}}\left[\frac{(x-a)^{\mu+1}+(b-x)^{\mu+1}}{(b-a)}\right]\end{aligned}$$

where $q>1$, $0\leq p\leq q$ *and*

$$K_2(1,m,t)=\begin{cases}\frac{M^{m}\left(\Gamma(\mu p+1)-\Gamma\left(\mu p+1,\ln M^{(m-1)}\right)\right)}{\left(\ln M^{(m-1)}\right)^{\mu p+1}} & ,M<1\\ \frac{1}{\mu p+1} & ,M=1\end{cases}.$$

Proof. If we set $\alpha=1$ in 2.5, the proof is completed. □

Corollary 5. *Let* $f:[0,\infty)\rightarrow(0,\infty)$ *be differentiable mapping with* $a,b\in[0,\infty)$ *such that* $a<b$. *If* $\left\vert f'(x)\right\vert^{q}$ *is logarithmically convex function with* $\left\vert f'(x)\right\vert^{q}\leq M, f'\in L[a,b]$ *and* $\mu>0$, *then the following inequality for fractional integrals holds:*

$$\begin{aligned}&\left\vert\frac{(x-a)^{\mu}+(b-x)^{\mu}}{b-a}f(x)-\frac{\Gamma(\mu+1)}{b-a}\left[J_{x^-}^{\mu}f(a)+J_{x^+}^{\mu}f(b)\right]\right\vert\\ =&\left(\frac{q-1}{\mu(q-p)+q-1}\right)^{\frac{q-1}{q}}\left(\frac{1}{\mu p+1}\right)^{\frac{1}{q}}\left[\frac{(x-a)^{2}+(b-x)^{2}}{(b-a)}\right]\end{aligned}$$

where $q>1$, $0\leq p\leq q$.

Proof. If we set $\alpha=m=1$ in 2.6, the proof is completed. □

Corollary 6. *Let* $f:[0,\infty)\rightarrow(0,\infty)$ *be differentiable mapping with* $a,b\in[0,\infty)$ *such that* $a<b$. *If* $\left\vert f'(x)\right\vert^{q}$ *is* $(\alpha,m)-$*logarithmically convex function with* $\left\vert f'(x)\right\vert^{q}\leq M, f'\in L[a,b]$ *and* $(\alpha,m)\in(0,1]\times(0,1]$, *then the following inequality*

holds:

$$\begin{aligned}&\left|f\left(x\right)-\frac{1}{b-a}\int_{a}^{b}f(u)du\right| \\ &\leq \left(\frac{q-1}{2q-p-1}\right)^{\frac{q-1}{q}}\left(K_{1}(\alpha,m,t)\right)^{\frac{1}{q}}\left[\frac{\left(x-a\right)^{2}+\left(b-x\right)^{2}}{\left(b-a\right)}\right]\end{aligned}\tag{2.7}$$

where $q>1$, $0\leq p\leq q$ *and*

$$K_{3}(\alpha,m,t)=\begin{cases}\frac{M^{m}\left(\Gamma(p+1)-\Gamma\left(p+1,\ln M^{\alpha(m-1)}\right)\right)}{\left(\ln M^{\alpha(m-1)}\right)^{p+1}} & ,M<1\\ \frac{1}{p+1} & ,M=1\end{cases}.$$

Proof. If we set $\mu=1$ in 2.5, the proof is completed. □

Corollary 7. *Let* $f:[0,\infty)\rightarrow(0,\infty)$ *be differentiable mapping with* $a,b\in[0,\infty)$ *such that* $a<b$. *If* $\left|f'\left(x\right)\right|^{q}$ *is* $\left(\alpha,m\right)-$*logarithmically convex function with* $\left|f'\left(x\right)\right|^{q}\leq M, f'\in L\left[a,b\right]$ *and* $(\alpha,m)\in(0,1]\times(0,1]$, *then the following inequality holds:*

$$\begin{aligned}&\left|f\left(x\right)-\frac{1}{b-a}\int_{a}^{b}f(u)du\right| \\ &\leq \left(\frac{1}{2}\right)^{\frac{q-1}{q}}\left(K_{1}(\alpha,m,t)\right)^{\frac{1}{q}}\left[\frac{\left(x-a\right)^{2}+\left(b-x\right)^{2}}{\left(b-a\right)}\right]\end{aligned}$$

where $q>1$, $0\leq p\leq q$ *and*

$$K_{4}(\alpha,m,t)=\begin{cases}\frac{M^{m}\left(\Gamma(2)-\Gamma\left(2,\ln M^{\alpha(m-1)}\right)\right)}{\left(\ln M^{\alpha(m-1)}\right)^{2}} & ,M<1\\ \frac{1}{2} & ,M=1\end{cases}.$$

Proof. If we set $p=1$ in 2.7, the proof is completed. □

REFERENCES

[1] R.-F. Bai, F. Qi and B.-Y. Xi, *Hermite-Hadamard type inequalities for the* $m-$ *and* $\left(\alpha,m\right)-$*logarithmically convex functions*, Filomat 27 (2013), 1-7.

[2] M.K. Bakula, J. Pečarić and M. Ribibić, *Companion inequalities to Jensen's inequality for* $m-$*convex and* $(\alpha,m)-$*convex functions*, J. Inequal. Pure and Appl. Math., 7 (5) (2006), Article 194.

[3] S.S. Dragomir and G. Toader, *Some inequalities for* $m-$*convex functions*, Studia University Babes Bolyai, Mathematica, 38 (1) (1993), 21-28.

[4] V.G. Miheşan, *A generalization of the convexity*, Seminar of Functional Equations, Approx. and Convex, Cluj-Napoca (Romania) (1993).

[5] G. Toader, *Some generalization of the convexity*, Proc. Colloq. Approx. *Opt.*, Cluj-Napoca, (1984), 329-338.

[6] G. Toader, *On a generalization of the convexity*, Mathematica, 30 (53) (1988), 83-87.

[7] A. Ostrowski, *Über die Absolutabweichung einer differentierbaren Funktion von ihren Integralmittelwert*, Comment. Math. Helv., 10, 226-227, (1938).

[8] M.E. Özdemir, H. Kavurmaci, E. Set, *Ostrowski's type inequalities for* $(\alpha,m)-$*convex functions*, KYUNGPOOK Math. J. 50 (2010) 371–378.

[9] H. Kavurmaci, M. Avci and M.E. Özdemir, *New Ostrowski type inequalities for* $m-$*convex functions and applications*, Hacettepe Journal of Mathematics and Statistics, Volume 40 (2) (2011), 135 – 145.

[10] M. Alomari and M. Darus, *Some Ostrowski type inequalities for convex functions with applications*, RGMIA Res. Rep. Coll., (2010) 13, 2, Article 3. [ONLINE: http://ajmaa.org/RGMIA/v13n2.php].

[11] E. Set, *New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals*, Comput. Math. Appl., 63 (2012) 1147-1154.

[12] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, J. Ineq. Pure and Appl. Math., 10(3), Art. 86 (2009).

[13] Z. Dahmani, *New inequalities in fractional integrals,* International Journal of Nonlinear Science, 9(4), 493-497 (2010).

[14] Z. Dahmani, *On Minkowski and Hermite-Hadamard integral inequalities via fractional integration*, Ann. Funct. Anal. 1(1), 51-58 (2010).

[15] Z. Dahmani, L. Tabharit and S. Taf, *Some fractional integral inequalities*, Nonl. Sci. Lett. A., 1(2), 155-160 (2010).

[16] M.Z. Sarıkaya, E. Set, H. Yaldiz and N. Başak, *Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities*, Mathematical and Computer Modelling, In Press.

[17] Z. Dahmani, L. Tabharit and S. Taf, *New generalizations of Grüss inequality using Riemann-Liouville fractional integrals*, Bull. Math. Anal. Appl., 2(3), 93-99 (2010).

[18] M.E. Özdemir, H. Kavurmacı and M. Avcı, *New inequalities of Ostrowski type for mappings whose derivatives are (α, m)-convex via fractional integrals*, RGMIA Research Report Collection, 15, Article 10, 8 pp (2012).

Ağri İbrahim Çeçen University, Faculty of Science and Letters, Department of Mathematics, Ağri, 04100, Turkey.

E-mail address: `ahmetakdemir@agri.edu.tr`